Dixie Animal Hospital, 14701 S. Dixie Hwy, Miami, FL, 33176 C. Tom Rainey, D.V.M. | Michael Pegelow, D.V.M. Phone #: 305-238-5161 | Fax #: 305-238-9297 dixieanimal@bellsouth.net | www.dixieanimalhospital.com



HOME CARE INSTRUCTIONS FOR SURGERY PATIENTS

Your pet has undergone anesthesia and surgery today. Your pet's ears were cleaned and nails were cut today at no charge. Please read the following recommendations to ensure a smooth recovery:

- 1. Your pet may be groggy due to the anesthesia and pain killers. Please use a leash, carry your pet or use a carrier to protect your pet when leaving the hospital.
- 2. When your pet arrives home, keep him/her quiet for a few hours. Offer water and feed a small amount at first to avoid vomiting. As the night goes on you may gradually offer more food. The next morning you may return to your pet's normal feeding schedule.
- 3. You may observe decreased activity or appetite for 1-2 days. However, please call us if your pet exhibits any of the following symptoms: refusal to drink water for longer than 1 day, loss of appetite for more than 2 days or excessive weakness, depression, vomiting or diarrhea. Some patients may not have a bowel movement for 1-3 days. Your pet may have a mild cough for a few days due to a breathing tube placed in the trachea.
- 4. Antibiotics are often prescribed if there is a possibility of infection; please finish all of the prescription as directed to treat and prevent infection. Pain killers/anti-inflammatories may also be prescribed for a few days.
- 5. Restrict your pet's activity for the next 7-14 days. Do not allow any strenuous activity (running, jumping, rough playing with other pets or children, etc...). Your pet should be confined indoors if possible and dogs should be leash walked only for a few days in order to restrict their activity and to monitor their bowel movements.
- 6. Your pet's incision should be monitored at least twice daily until healed. A small amount of bloody seepage is normal the night of the surgery. If you notice any continued drainage or pus or any excessive redness or swelling please call us. You may clean the incision with warm water and a soft paper towel. You may apply triple antibiotic ointment/Neosporin as needed to the incision. You may use 1% hydrocortisone ointment on the shaved skin around the incision as needed to treat irritation caused by the clipping of the hair.
- 7. If your pet has sutures at the incision you will need to bring him/her back in 10-14 days for us to remove the sutures and examine the incision. Please call for an appointment for suture removal. Please do not bathe your pet until his/her follow up appointment unless it is absolutely necessary.
- 8. If your pet is excessively licking at his/her incision you will need to contact us about getting a special "cone" collar that prevents your pet from licking at the incision. These collars are also available at Petsmart and Petco.
- 9. If your pet was spayed or neutered you should feed 10-20% less food on a daily basis to avoid excessive weight gain. Once healed, exercise has many benefits to a long, healthy life.

Please call (305) 238-5161 if you have any questions or concerns.